

KINEMATICS AND EXCITATION OF THE RAM PRESSURE STRIPPED IONIZED GAS FILAMENTS IN THE COMA CLUSTER OF GALAXIES¹

MICHITOSHI YOSHIDA², MASAFUMI YAGI³, YUTAKA KOMIYAMA³, HISANORI FURUSAWA⁴, NOBUNARI KASHIKAWA³, TAKASHI HATTORI⁵ AND SADANORI OKAMURA⁶

Draft version February 7, 2012

ABSTRACT

We present the results of deep imaging and spectroscopic observations of very extended ionized gas (EIG) around four member galaxies of the Coma cluster of galaxies: RB199, IC 4040, GMP 2923 and GMP 3071. The EIGs were serendipitously found in an H α narrow band imaging survey of the central region of the Coma cluster. The relative radial velocities of the EIGs with respect to the systemic velocities of the parent galaxies from which they emanate increase almost monotonically with the distance from the nucleus of the respective galaxies, reaching $\sim -400 - -800$ km s⁻¹ at around 40 – 80 kpc from the galaxies. The one-sided morphologies and the velocity fields of the EIGs are consistent with the predictions of numerical simulations of ram pressure stripping. We found a very low-velocity filament ($v_{\text{rel}} \sim -1300$ km s⁻¹) at the southeastern edge of the disk of IC 4040. Some bright compact knots in the EIGs of RB199 and IC 4040 exhibit blue continuum and strong H α emission. The equivalent widths of the H α emission exceed 200 Å and are greater than 1000 Å for some knots. The emission line intensity ratios of the knots are basically consistent with those of sub-solar abundance H II regions. These facts indicate that intensive star formation occurs in the knots. Some filaments, including the low velocity filament of the IC 4040 EIG, exhibit shock-like emission line spectra, suggesting that shock heating plays an important role in ionization and excitation of the EIGs.

Subject headings: galaxies: clusters: individual (Coma) — galaxies: clusters: intracluster medium — galaxies: evolution — galaxies: kinematics and dynamics — intergalactic medium

1. INTRODUCTION

Clusters of galaxies are ideal laboratories for investigating environmental effects in galaxy evolution. The extreme environments of the clusters — the high number density of galaxies, strong gravitational field, and dense hot gas filling inter-galactic space — allow us to study how these environmental factors affect the evolution of cluster member galaxies. A substantial body of observational evidence suggests that cluster environment significantly affects cluster galaxy evolution; for example, the morphology - density relation (Dressler 1980; Postman & Geller 1984; Dressler 1994; van der Wel et al. 2007), the redshift - blue galaxy fraction relation (Butcher & Oemler 1978, 1984), or the density - H I deficiency relation (Cayatte et al. 1990, 1994; Solanes et al. 2001). Researchers have proposed various physical mechanisms potentially responsible for these observational facts (Boselli & Gavazzi 2006), including galaxy-galaxy interaction (Byrd & Valtonen 1990), galaxy harassment (Moore et al. 1996), tidal interaction between galaxies and cluster potential (Byrd & Valtonen

1990; Henriksen & Byrd 1996), galaxy starvation (Bekki & Couch 2003), and ram pressure stripping (Gunn & Gott 1972, hereafter referred as to RPS).

Although it is not clear which mechanism plays the most important role in the morphological and color evolution of cluster galaxies, it is at least certain that rapid removal of the interstellar gas of galaxies is a key factor (Okamoto & Nagashima 2003; van Gorkom 2004; Boselli et al. 2008). In particular, RPS caused by fast interaction between galaxies and the hot intra-cluster medium (ICM) must play a very important role in gas removal (Vollmer et al. 2001b; Fujita 2004; Cortese et al. 2011). A number of observational studies have focused on this subject. The H I gas of galaxies is systematically deficient in the cores of nearby rich clusters (Cayatte et al. 1990, 1994; Bravo-Alfaro et al. 2000, 2001). Deep H I imaging revealed one-sided H I gas flows around late-type galaxies in the Virgo cluster (Oosterloo & van Gorkom 2005; Chung et al. 2007, 2009). Asymmetric distributions of molecular gas found in some cluster galaxies also suggest that the molecular gas is affected by ram pressure from the hot ICM (Vollmer et al. 2001a; Vollmer & Huchtmeier 2003; Vollmer et al. 2008, 2009; Sivanandam, Rieke & Rieke 2010). Recently, elongated X-ray gas-flows from galaxies have been found in nearby clusters (Sun, Donahue & Voit 2007; Sun et al. 2010; Randall et al. 2008). In addition, many case studies have focused on ionized gas flows from cluster member galaxies probed by H α emission (Kenney & Koopmann 1999; Gavazzi et al. 2001, 2003; Yoshida et al. 2002, 2004; Sakai et al. 2002; Chemin et al. 2005; Cortese et al. 2006; Kenney et al. 2008). Some pieces of evidence of

¹ Based on data collected at the Subaru Telescope, which is operated by the National Astronomical Observatory of Japan.

² Hiroshima Astrophysical Science Center, Hiroshima University, Higashi-Hiroshima, Hiroshima 739-8526, Japan; yoshidam@hiroshima-u.ac.jp

³ Optical and Infrared Astronomy Division, National Astronomical Observatory, Mitaka, Tokyo 181-8588, Japan.

⁴ Astronomical Data Center, National Astronomical Observatory of Japan, Mitaka Tokyo 181-8588, Japan.

⁵ Subaru Telescope, National Astronomical Observatory of Japan, 650 North A'Ohoku Place, Hilo, HI 96720, USA.

⁶ Department of Astronomy, University of Tokyo, Tokyo 113-0033, Japan.

dust stripping in the Virgo galaxies have also been reported (Crowl et al. 2005; Abramson et al. 2011).

We serendipitously discovered a number of very extended ionized gas filaments (EIG) around the member galaxies of the Coma cluster (Yagi et al. 2007; Yoshida et al. 2008, hereafter YY08; Yagi et al. 2010, hereafter YY10). The filaments are extended toward one side of the parent galaxies (YY10). The most remarkable example of the one-sided extension is the 60-kpc straight $H\alpha$ filament found around D100 (Yagi et al. 2007). Another interesting object is the complex of ionized gas filaments, star-forming blue knots and young stellar filaments extended from an E+A galaxy RB199 (“fireballs”; YY08). The morphology of the latter very much resembles strings of compact blue knots around the two galaxies falling into $z \approx 0.2$ clusters found by Cortese et al. (2007). Cortese et al. (2007) interpreted these peculiar features as star-forming regions in the gas stripped from the parent galaxies by the combined action of tidal interaction and RPS. The fireballs is a low-redshift counterpart of these features (YY08). Recently, Hester et al. (2010) reported a similar structure around a low surface brightness galaxy IC 3418 in Virgo cluster (see also Fumagalli et al. 2011).

YY10 found some properties specific to the EIGs’ parent galaxies. The radial velocities of the parent galaxies are distributed either at the blue or at the red edge of the velocity distribution of Coma member galaxies. In addition, almost all the parent galaxies have blue colors; about 60% of the parent galaxies are distributed well below (bluer side) from the color-magnitude relation (red sequence) of the early type galaxies of the Coma cluster. These facts suggest that the parent galaxies are late-type galaxies which have recently fallen into the cluster. They have not yet been well processed in the cluster environment. Further, YY10 classified the EIGs into three categories: (1) disk star-formation + connected $H\alpha$ gas, (2) no disk star-formation with connected $H\alpha$ gas, and (3) detached $H\alpha$ gas.

Smith et al. (2010) found on *GALEX* images one-sided UV tails around a number of Coma member galaxies whose morphologies resemble the EIGs. More than half of their objects overlapped the EIG galaxies discovered by YY10. They assessed the statistical characteristics of these UV tail galaxies and concluded that these galaxies are newcomers to the Coma cluster (Smith et al. 2010). Good correlation between the EIGs and the UV tails suggests that these two kinds of features were created by the same mechanism; the most plausible one is RPS. Furthermore, this indicates active star formation occurs in the stripped gas in the cluster environment.

Here we present the results of optical spectroscopic observations of the EIGs around four Coma member galaxies. We assumed that the cosmological parameters (h_0 , Ω_m , Ω_λ) = (0.73, 0.24, 0.72) and the distance modulus of the Coma cluster is 35.05 (Yagi et al. 2007). The linear scale at the Coma cluster is 474 pc arcsec $^{-1}$ under this assumption.

2. OBSERVATION AND DATA REDUCTION

Deep spectroscopic observations of four Coma member galaxies, RB199, IC 4040 (GMP 2559), GMP 2923 and GMP 3071 was performed using the faint object spectrograph FOCAS (Kashikawa et al. 2002) attached to the

Subaru Telescope on 20 May 2009. Two multi-slit masks for each of RB199 and IC 4040 were used to cover the complicated structure of the emission line regions over as wide an area as possible. GMP 2923 and GMP 3071 were observed using one multi-slit mask. The positions and lengths of the slits are shown in Figures 1 – 4. The width of each small slit was 0".8. The lengths of the slits ranged from 10" to 30". A 300 grooves mm $^{-1}$ grism, whose straight light travel wavelength is 6500 Å, was used. The full width at half maximum (FWHM) of the sky emission lines around 6700 Å is 7.8 Å, which means that the spectral resolving power R of our spectroscopy is 850 (the velocity resolution is about 350 km s $^{-1}$) over $H\alpha$ region. Table 1 presents the observation log.




FIG. 1.— Slit positions of the spectroscopy of the extended ionized gas (EIG) of RB199 (“fireballs”: YY08) overlaid on a composite image of the $H\alpha$ narrow band and B band images. All the images used in Figures 1 – 4 were taken with Suprime Cam attached to Subaru Telescope. Details of the observations of these images were described previously in YY10.

TABLE 1
SPECTROSCOPY OBSERVATION LOG

mask ID	object	center	PA	exposure
MS0631	IC 4040	$13^h00^m43.2^s +28^\circ03'06''$	215°	$6 \times 600\text{sec}$
MS0632	IC 4040	$13^h00^m38.4^s +28^\circ04'42''$	125°	$3 \times 300\text{sec} + 3 \times 600\text{sec}$
MS0632-off	IC 4040	$13^h00^m38.6^s +28^\circ04'45''$	125°	$1 \times 600\text{sec}$
MS0634	RB199	$12^h58^m43.3^s +27^\circ46'19''$	150°	$6 \times 600\text{sec}$
MS0635	RB199	$12^h58^m38.7^s +27^\circ44'35''$	210°	$6 \times 600\text{sec}$
MS0636	GMP2923,3071	$12^h59^m58.4^s +27^\circ44'58''$	60°	$2 \times 600\text{sec}$




FIG. 2.— Slit positions of the spectroscopy of the EIG of IC 4040 overlaid on the net $H\alpha$ image.

Data reduction was performed in standard manner. The bias level of each frame was estimated with average count of the over-scan region of the CCD and then subtracted from each frame. Flat-fielding was done using dome flat frames. Wavelength calibration was performed using sky emission lines. The rms residual of wavelength




FIG. 3.— Slit positions of the spectroscopy of the EIG of GMP 2923 overlaid on the $H\alpha$ narrow band image.




FIG. 4.— Slit positions of the spectroscopy of the EIG of GMP 3071 overlaid on the $H\alpha$ narrow band image.

fitting for sky lines was $0.14 \text{ \AA}$ in the wavelength region of $6000 \text{ \AA} - 7000 \text{ \AA}$. Because there are no strong sky emission lines blueward of $5500 \text{ \AA}$, wavelength calibration is not accurate as in the longer wavelength region. The rms error was $0.3 \text{ \AA}$ at $5300 \text{ \AA}$. Thus, we measured the radial velocities of the EIGs using the $H\alpha$ line only.

Flux calibration was initially conducted using spectrophotometric standard star data. However, the standard star data were taken with an order sorting filter (Y47) whose cut-off wavelength was $4700 \text{ \AA}$, while the filter was not used for the observation of the EIGs. Thus, we had to correct the transmission curve of Y47 to ensure accurate flux calibration for the wavelength region blueward of $4700 \text{ \AA}$. We divided the initial flux-calibrated frames by the transmission curve of Y47, which was normalized to 1 at the maximum transmission wavelength ($6000 \text{ \AA}$). We checked validity of the above recalibration procedure by comparing the calibrated spectrum of an elliptical galaxy (SDSS J125845.26+274655.0) taken with one slit of the mask MS0635 with the spectra of eight early type galaxies listed in Kennicutt (1998). The type of the reference galaxies ranges from E0 to S0. All the

spectra were normalized at 6000 Å. The rms relative difference between the spectrum of our E galaxy and the spectrum of each reference early type galaxy was less than 7% for the wavelength region between 4300 Å and 7000 Å. Thus we adopted that the recalibration we made was correct enough for further analysis. The signal-to-noise ratio of the measured spectra shorter than 4300 Å was too low to enable confident comparison, so we do not include spectra shorter than 4300 Å in the following analysis and discussion.

We fitted gaussian functions to the emission lines to measure the fluxes and the velocities of the lines. Although it was difficult to measure velocity dispersion accurately because of low spectral resolution of our spectra, we obtained rough estimation of the velocity dispersion σ by applying a simple decomposition scheme; $\sigma = \sqrt{\sigma_{\text{obs}}^2 - \sigma_{\text{sky}}^2}$, where σ_{obs} and σ_{sky} are the dispersions of the gaussian functions fitted to the observed emission line and that of the sky line, respectively ($f(\lambda) \propto \exp(-\lambda^2/2\sigma^2)$; $\sigma = \text{FWHM}/2\sqrt{2\ln 2}$). One spectral resolution unit was sampled with eight pixels of the CCD. This over-sampling enabled us to measure the emission line widths with an accuracy at 10% of the FWHM of the sky lines. Since the FWHM of the sky lines was 7.8 Å, we resolved the emission lines whose FWHMs were wider than 8.6 Å, which means that the lower limit of σ we could measure was $\approx 70 \text{ km s}^{-1}$ at 6700 Å.

Imaging data of the target galaxies in deep narrow-band H α and broad-band B and R_C bands used in this paper are those presented in YY10 for morphological study of the EIGs around these galaxies. They were taken using Suprime Cam (Miyazaki et al. 2002) attached to the Subaru Telescope. The data were collected over three observing runs from 2006 - 2009. See YY10 for the details of the imaging observations and data reduction.

3. RESULTS

3.1. The morphologies and kinematics of the EIGs

The overall morphological characteristics of the EIGs of our target galaxies were previously reported by YY10. Here, we summarize the details of the morphologies and give an interpretation to them based on the newly obtained kinematical information.

3.1.1. RB199

The B , R_C and H α images of RB199 (fireball) were shown in our previous work (YY08). The characteristics of the EIG of RB199 can be summarized as follows: (1) The EIG consists of a number of bright star-forming knots, elongated blue filaments and more extended H α emitting gas. Total length of the EIG is about 80 kpc. (2) The absolute magnitudes of the bright knots range from $-12 - -13$ mag, which are comparable to nearby dwarf galaxies. (3) The bright knots emit both blue continuum light and H α emission. The H α emitting regions are offset from the peak of the blue continuum emission. (4) The blue filaments are connected to the main body of the galaxy, while the H α filaments are connected to the tips of the blue filaments. (5) The bright knots and blue filaments are detected in UV light by the *GALEX* satellite.




FIG. 5.— The velocity field of the EIG of RB199. The systemic velocity of the galaxy is determined by the H α absorption line of the nucleus.




FIG. 6.— The velocity fields of the EIG of RB199 and the low velocity filaments overlapping H α filament 2.

Figure 5 shows the velocity field of the EIG of RB199. Relative radial velocities of the EIG filaments with respect to the systemic velocity of RB199 increase with the distance from the nucleus at almost a constant rate of $-7 \text{ km s}^{-1} \text{ kpc}^{-1}$. The relative radial velocity of the most distant cloud of the EIG, H α cloud 2 (YY08), is roughly consistent with the extrapolation of the line of constant increase, which suggests that this cloud is really a part of the EIG.

We found that a faint H α component with a velocity of $\approx 6250 \text{ km s}^{-1}$ overlaps along the line of sight to H α cloud 2. Figure 6 shows the velocity field of the RB199 EIG with this slow velocity (SV) component. The velocity of this SV component is slightly lower than that of the mean velocity ($\approx 6900 \text{ km s}^{-1}$) of the Coma cluster. The velocity difference between the SV component and H α cloud 2 is $\sim 2000 \text{ km s}^{-1}$, suggesting that the SV component is not associated with the RB199 EIG.

3.1.2. IC 4040

Figure 7 presents a pseudo-color image (blue, green and red colors represent B , R_C , and H α images, respectively) of the spiral galaxy IC 4040 taken with Subaru Suprime-Cam (YY10, reproduced by permission of the American Astronomical Society; AAS). The figure clearly shows a very extended EIG expelled from the galaxy disk.




FIG. 7.— A pseudo color image (B , R_C and $H\alpha$ composite: left panel) and net $H\alpha$ image (right panel) of IC 4040. This figure is reproduced from YY10 by permission of the AAS.

The structure of the EIG is very complicated. Two major streams extend toward the southeast and south from the galaxy. The southeast stream of the EIG is elongated almost linearly along the major axis of the galaxy and its root is connected to the nuclear star-forming region. This southeast stream reaches ~ 50 kpc from the nucleus. Figure 8 shows the detailed structure of the central bright part of the EIG. The nucleus of the galaxy is surrounded by several bright star-forming knots. This complex of the star-forming regions is truncated at the northwestern side of the nucleus. The shape of the ionized gas distribution indicates a large-scale bow shock and a giant gas flow toward the southeast from the nucleus. The opening angle of the gas flow is about 65° and almost symmetric with respect to the major axis of the galaxy. Several bright $H\alpha$ knots (NW-K1, SE-K1, etc. see Figure 8) surround the nucleus. The brightest one, SE-K4, has a size of $r \sim 0.5$ kpc and is located at 7 kpc from the nucleus. A large complex of diffuse $H\alpha$ clouds/filaments appears at around 15 kpc from the nucleus. Further away, a tangle of several filaments, with typical width of 1 kpc, extends out to 50 kpc. For convenience, we refer to the large complex and the filaments between 20 kpc and 30 kpc as SE-F1 and SE-F2, respectively. As described later, our spectroscopic observations revealed that the SE-F1 consists of at least two kinematically-different components, a low-velocity (LV) component and a high-velocity (HV) component.

The $B - R_C$ color image shown in Figure 9 reveals that the southeast stream is very dusty. The close spatial correlation between the dust and the EIG and the one-sided morphology of the dust suggest that the ionized gas and the dust are stripped together and are mixed with each




FIG. 8.— Details of the morphology of the $H\alpha$ emitting gas of IC 4040 around the galaxy disk and nucleus. In the central region of the galaxy, several bright $H\alpha$ knots and complicated structure of ionized gas filaments/clouds can be seen. The $H\alpha$ gas is truncated at the north-western side and exhibits bow shock like morphology.




FIG. 9.— $B - R_C$ color image (gray-scale) of IC4040. The darker color represents a larger $B - R_C$. Contour map of the $H\alpha$ gas distribution is overlaid on the $B - R_C$ image.

other in the stripped flow. Both the dust stream and the southeast part of the EIG extend in the same direction as the H I gas tail found by Bravo-Alfaro et al. (2000, 2001). Recent far-infrared/submm survey of the Virgo cluster with *Herschel* revealed that dust disks are truncated in H I deficient galaxies (Cortese et al. 2010). These results suggest that dust in galaxies can be stripped in the cluster environment. The dust stream in IC 4040 exemplifies a dust stripping site observed in a cluster.

The EIG is widely distributed south of IC 4040. The southern part of the EIG consists of three conspicuous bright knots (S-K1, S-K2 and S-K3; see Figure 2), several giant complexes of clouds/filaments (S-F1, S-F2, etc.) and many small filaments.

The three bright knots are blue in color. These features have been detected in the UV light with *GALEX* (Smith et al. 2010). Of the three knots, knot 2 is the

brightest. Smith et al. (2010) observed a Wolf-Rayet feature in their Keck spectrum, suggesting this knot is very young (younger than 5 Myr). Elongated H α filaments are connected to these knots. All the connected filaments have head-tail morphologies whose position angles are the same as the southeast stream.

Many ionized gas clouds/filaments are distributed over $30 \text{ kpc} \times 80 \text{ kpc}$ in the south area of IC 4040. The brightest one is a thick loop-like filament at about 40 kpc from the galaxy (S-F1 in Figure 2). S-F2 is another bright filament connected to S-F1. S-F3 and S-F4 are diffuse filaments seen near an early type galaxy GMP 2551 (Figure 7). Other than these filaments, many small filaments are seen between S-F2 and IC 4040.




FIG. 10.— The velocity field of the EIG of IC 4040. Green dots, blue dots and red dots represent the data points of the slit masks of MS0631, MS0632, and MS0632-off, respectively.

Figure 10 shows the velocity field of the EIG of IC 4040. The brightness of the EIG of IC 4040 allowed us to trace the velocity field along each slit. The IC 4040 EIG has a velocity field similar to that of the RB199 EIG; the relative radial velocities of the filaments increase almost monotonically with the distance from the nucleus (Figure 10). Within 5 kpc of the nucleus, the H α gas along the galaxy major axis follows the galaxy rotation. The acceleration of the EIG begins at 5 kpc from the nucleus. The overall acceleration rate in a range of 5 – 80 kpc is $\sim -10 \text{ km s}^{-1} \text{ kpc}^{-1}$.

We noted several peculiar features in the velocity field of the IC 4040 EIG. The most remarkable one is a very low-velocity component at $\sim 12 \text{ kpc}$ from the nucleus. This component is spatially associated with the SE-F1 (see Figure 2); we call it SE-F1 (LV). The relative velocity of the SE-F1 (LV) is -1300 km s^{-1} with respect to the systemic velocity of the galaxy. Its velocity dispersion reaches $\approx 800 \text{ km s}^{-1}$ (Figure 11). In contrast, the southwest side of the SE-F1 is much faster than the SE-F1 (LV) and almost follows the overall motion of the EIG. This fast component, called SE-F1 (HV) has a relative velocity of $\approx -300 \text{ km s}^{-1}$ and a velocity dispersion of $\approx 400 \text{ km s}^{-1}$. Figure 12 shows the two-dimensional spectra of the circumnuclear H α gas and the southeast filaments. In this figure, large velocity shift and wide velocity dispersion of SE-F1 (LV) are clearly seen.

The S-F1 has a large velocity gradient ($\sim 70 \text{ km s}^{-1} \text{ kpc}^{-1}$) within it. The filament has a relative




FIG. 11.— The velocity dispersion of the EIG of IC 4040. The color difference is the same as Figure 10.




FIG. 12.— Two dimensional spectra of the SE filaments and the nuclear H II regions of IC 4040. Middle panel shows a pseudo color image of the H α emitting filaments.

velocity of $\approx -700 \text{ km s}^{-1}$ at the edge of the galaxy. At the opposite edge, the relative velocity reaches -450 km s^{-1} . The S-F3 and S-F4 are highly turbulent and the velocity dispersions are $\sim 200 - 400 \text{ km s}^{-1}$.

3.1.3. GMP 2923 and GMP 3071

The EIG around GMP 2923 is shown in Figure 4d in YY10. Two galaxies — GMP 2923 and GMP 2945 — overlap in the line of sight near the EIG (Figure 3). We measured the radial velocity of the early-type galaxy GMP 2945 with the same slit mask as that used for the EIG; it is 6500 km s^{-1} , much slower than those of the EIG and GMP 2923. Thus GMP 2945 is kinematically independent from the EIG.

The EIG of GMP 2923 is characterized by two knotty straight streams (see Figure 4d in YY10). The opening angle of the streams is ≈ 20 degrees. The root of the streams is detached from the galaxy. The EIG is extended to $\sim 50 \text{ kpc}$ from GMP 2923.

The EIG of GMP 3071 consists of slightly wiggled narrow filaments (Figure 4f in YY10). The root of the EIG is connected to the nucleus of the galaxy. The angle between the galaxy major axis and the EIG is ≈ 40 degrees and the length of the EIG is $\sim 25 \text{ kpc}$.




FIG. 13.— The velocity field of the EIG of GMP 2923.




FIG. 14.— The velocity field of the EIG of GMP 3071.

The kinematics of both the GMP 2923 and GMP 3071 EIGs also reveal a monotonically-increasing pattern. Figure 13 and 14 show the velocity fields of the EIGs of the two galaxies. The acceleration rates are $\approx -10 \text{ km s}^{-1} \text{ kpc}^{-1}$ and $\approx -25 \text{ km s}^{-1} \text{ kpc}^{-1}$ for GMP 2923 and GMP 3071, respectively.

The kinematics of the EIG of GMP 3071 is rather complex in the galaxy disk, showing a rapid drop-and-rise whose amplitude is $\approx 200 \text{ km s}^{-1}$ (Figure 14). The net $\text{H}\alpha$ image taken by YY10 revealed that the ionized gas near the nucleus has a wiggled chain of small blobs. Together with the kinematics, this may suggest a helical motion of the ionized gas caused by the stripping force coupled with galactic rotation.

3.2. Excitation of the EIGs

Most of the EIG extended filaments are too faint to detect emission lines other than $\text{H}\alpha$. We could detect diagnostic emission lines such as $[\text{S II}]$ or $[\text{N II}]$ only in bright knots and some bright filaments in the EIGs of RB199 and IC 4040. Table 2 summarizes the measured fluxes of the emission lines for bright parts of the EIGs of RB199 and IC 4040.

Figure 15 shows the spectra of the bright knots and the nucleus of RB199. The bright knots of RB199 are characterized by strong emission lines and blue continuum. The equivalent widths of $\text{H}\alpha$ are $310 \text{ \AA}$, $230 \text{ \AA}$,




FIG. 15.— Spectra of the bright knots of the EIG and the nucleus of RB199.

and $1500 \text{ \AA}$ for knot 1, 2, and 5, respectively. The most interesting feature of the bright knots spectra is the weak $[\text{N II}]6548/6583$ emission. In knot 2 or knot 3, $[\text{S II}]6717/6731$ lines are stronger than $[\text{N II}]$. In knot 3, even $[\text{O I}]6300$ is stronger than $[\text{N II}]$. In knot 1 and knot 5, very weak $[\text{N II}]$, $[\text{S II}]$ and $[\text{O I}]$ and relatively strong $[\text{O III}]5007$ indicate that these knots are metal-deficient H II regions. The spectra of knot 2 and knot 3 reveal strong $[\text{S II}]$ and $[\text{O I}]$ lines, suggesting that shock heating contributes the gas excitation. The nucleus of RB199 has a spectrum typical of post-starburst galaxies. It is characterized by strong Balmer absorption lines, very weak metal-absorption lines, and a blue continuum.

Figure 16 shows the spectra of the bright knots and filaments of the EIG of IC 4040 with the nuclear spectrum of IC 4040 itself. The spectrum of S-K2 of the IC 4040 EIG is that of a typical metal-poor star-forming region. The $\text{H}\alpha$ line is very strong in S-K2; the equivalent width reaches $\sim 1000 \text{ \AA}$. In the blue region of the spectrum, we confirmed a weak Wolf-Rayet feature (Figure 17) which was found by Smith et al. (2010). S-K3 shows relatively strong $[\text{N II}]$ and $[\text{S II}]$ suggesting shock excitation may play a role in the gas excitation. The spectra of the nucleus and S-K4 resemble each other, while the continuum of S-K1 is redder than that of the nucleus. It reflects the dusty gas flow in the disk of the galaxy (Figure 9) and

TABLE 2
EMISSION LINE FLUXES AND INTENSITY RATIOS OF THE EIGs.

ID	distance ^a	length ^b	$f_{\text{H}\alpha}^c$	$f_{\text{H}\beta}^d$	[NII]/H α^e	[SII]/H α^e	[OI]/H α^e	[OIII]/H β^e
IC 4040								
nucleus	—	0.59	271	85.3	-0.47 ± 0.01	-0.52 ± 0.01	-1.60 ± 0.01	-0.36 ± 0.01
NW-K1	1.2	0.44	221	51.2	-0.48 ± 0.01	-0.61 ± 0.01	-1.69 ± 0.02	-0.54 ± 0.01
SE-K1	1.1	0.74	303	53.6	-0.32 ± 0.01	-0.57 ± 0.01	-1.36 ± 0.01	-0.43 ± 0.01
SE-K2	3.8	0.90	47.1	17.1	-0.40 ± 0.01	-0.44 ± 0.01	-1.22 ± 0.03	-0.51 ± 0.03
SE-K3	2.5	0.74	107	23.9	-0.43 ± 0.01	-0.58 ± 0.01	-1.42 ± 0.02	-0.27 ± 0.01
SE-K4	4.8	0.74	194	37.1	-0.51 ± 0.01	-0.61 ± 0.01	-1.61 ± 0.02	-0.17 ± 0.01
S-K1	12.3	0.59	2.1	—	-0.49 ± 0.10	-0.12 ± 0.07	-0.48 ± 0.10	—
S-K2	17.0	0.74	172	59.1	-1.04 ± 0.01	-1.05 ± 0.01	-1.98 ± 0.04	0.50 ± 0.01
S-K3	21.7	0.74	14.6	5.3	-0.37 ± 0.02	-0.40 ± 0.02	-1.41 ± 0.12	-0.05 ± 0.04
S-K4	47.9	0.74	3.2	—	—	—	—	—
SE-F1(LV)	12.8	2.22	16.4	3.9	-0.12 ± 0.03	-0.51 ± 0.09	-0.64 ± 0.09	0.04 ± 0.13
SE-F1(HV)	13.8	3.11	28.4	3.4	-0.16 ± 0.03	-0.38 ± 0.06	-0.63 ± 0.06	0.12 ± 0.21
SE-F2	26.7	1.78	14.8	—	—	—	—	—
S-F1N	37.8	0.59	5.7	2.1	-0.44 ± 0.04	-0.27 ± 0.04	-0.77 ± 0.07	-0.42 ± 0.15
S-F1S	39.2	0.74	5.5	2.1	-0.47 ± 0.05	-0.42 ± 0.06	-0.90 ± 0.11	-0.36 ± 0.18
S-F2	40.9	1.04	2.8	—	-0.25 ± 0.08	-0.27 ± 0.10	—	—
S-F3	68.8	2.37	6.3	—	-1.08 ± 0.22	—	—	—
RB199								
knot1	22.8	1.18	23.5	8	-1.02 ± 0.03	-0.67 ± 0.02	-1.52 ± 0.11	0.19 ± 0.02
knot1e	23.7	1.04	3.8	0.7	-1.06 ± 0.33	-0.24 ± 0.09	-0.61 ± 0.14	-0.25 ± 0.44
knot2	24.1	1.48	37.9	13	-0.86 ± 0.02	-0.29 ± 0.01	-1.06 ± 0.03	0.07 ± 0.02
knot3	34.8	0.89	4.9	2	-0.92 ± 0.09	-0.26 ± 0.03	-0.99 ± 0.10	-0.22 ± 0.11
knot4	36.2	0.74	1.9	—	—	—	—	—
knot5	34.9	1.04	21.6	8	-1.14 ± 0.06	-0.65 ± 0.03	-1.45 ± 0.12	0.37 ± 0.02
H α filament 1(N)	43.2	2.51	1.3	—	—	—	—	—
H α filament 1(S)	50.4	1.18	2.4	—	—	—	—	—
H α filament 2	39.8	0.89	2.7	—	—	—	—	—
H α cloud 2	76.3	1.48	1.9	—	—	—	—	—

(a) Distance from the nucleus of the parent galaxy in unit of kpc.

(b) Length along the slit in unit of kpc.

(c) H α flux in unit of 10^{-17} ergs s $^{-1}$ cm $^{-2}$.

(d) H β flux in unit of 10^{-17} ergs s $^{-1}$ cm $^{-2}$.

(e) Logarithmic value of the emission line intensity ratio.

S-K4 is embedded in the dusty flow.

Certain low ionization forbidden lines, [S II], [O I] and [N II], are enhanced relative to H α in some extended filaments of the IC 4040 EIG: SE-F1, S-F1 and S-F3 (Figure 16). An extended filament attached to the tip of knot 1 (east edge of knot 1) of RB199 also exhibits enhanced [S II] and [O I], as shown in the bottom panel of Figure 15. Other extended filaments are too faint to measure the intensities of emission lines other than H α .

Figures 18, 19 and 20 are emission line diagnostic diagrams (Veilleux & Osterbrock 1987) of the EIG knots and filaments of RB199 and IC 4040. These diagrams clearly show the above-mentioned characteristics of the EIG spectra. The relative weakness of [N II]6583 in the RB199 EIG is conspicuous in Figure 18; the emission line intensity ratio $\log([\text{N II}]/\text{H}\alpha) \leq -1.0$ for the knots of the RB199 EIG. Extragalactic H II regions, which have low [N II]/H α ratios, also generally have low [S II]/H α and [O I]/H α ratios. Figure 21 shows the abnormally weak [N II] emission of the RB199 knots. The data points of knots 1e, 2 and 3 are well displaced from the main sequence of star-forming galaxies, whereas knots 1 and 5 are embedded in the sequence.

The emission line ratios of the bright knots and the nucleus of IC 4040 with the exception of S-K1 are con-

sistent with those of star-forming galaxies (Figures 18, 19, 20 and 21). Among these, it is clear that S-K2 of IC 4040 shows an emission line spectrum like blue compact dwarf galaxies. In contrast, some extended filaments have LINER-like spectra. In particular, the [O I] line is enhanced in these features (Figure 20). This [O I] enhancement suggests that shock heating plays an important role in ionization and excitation of these features. The [N II] weak features in the RB199 EIG also reveal similar [O I] enhancement, suggesting that shock heating is important for these peculiar features. However, the abnormally weak [N II] line of these features cannot be explained by ordinary shock heating models. Low velocity shock heated gas which enhances [O I] and [S II] overlapped on low metal abundance H II region might produce such peculiar emission-line intensity ratios.

We tried to estimate the metal abundance of the bright knots around RB199 and IC 4040 using the model calculations by Kewley & Dopita (2002). We used [N II]/[O III] and [N II]/H α ratios to estimate the metal abundance. The kinematical characteristics and oxygen abundances of the bright knots are shown in Table 3. The emission-line widths of most of the bright knots except for S-K2 are too narrow to resolve with our spectral resolution, indicating that the knots are much colder

TABLE 3
KINEMATICS AND ABUNDANCES OF THE BRIGHT KNOTS.

ID	v_{rel} (km s $^{-1}$)	σ^a (km s $^{-1}$)	$\log(\text{O}/\text{H})+12$	$Z_{\odot}^b$
RB199				
knot1	-104 ± 16	< 70	8.38 ± 0.05	0.49 ± 0.08
knot2	-194 ± 15	< 70	8.48 ± 0.05	0.62 ± 0.10
knot3	-210 ± 17	< 70	8.40 ± 0.03	0.51 ± 0.07
knot5	-240 ± 15	< 70	8.44 ± 0.05	0.56 ± 0.09
IC 4040				
S-K2	-241 ± 18	93 ± 20	8.59 ± 0.03	0.79 ± 0.10
S-K3	-268 ± 16	< 70	8.88 ± 0.05	1.55 ± 0.25
SE-K4	-210 ± 17	< 70	8.40 ± 0.02	0.51 ± 0.06

(a) Velocity dispersion σ of H α line derived by gaussian fitting ($f(\lambda) \propto \exp(-\lambda^2/2\sigma^2)$).

(b) Oxygen abundances relative to the solar value derived by Asplund et al. (2009) ($\log(\text{O}/\text{H})+12 = 8.69 \pm 0.05$).




FIG. 16.— Spectra of the bright knots and filaments of the EIG and the nucleus of IC 4040.

than the extended filaments. As suggested by the emission line diagnostic diagrams (Figures 18, 19 and 20), the bright knots except for S-K3 and SE-K4 have basically low oxygen abundances; $\log(\text{O}/\text{H})+12 \sim 8.4 - 8.6$. These values correspond to $0.5 - 0.8 Z_{\odot}$. S-K3 of IC 4040 shows super solar abundance ($\sim 1.6 Z_{\odot}$). Because this knot is located at marginal region between H II regions and LINERs in the diagnostic diagrams, shock heating may contribute enhancement of the forbidden lines, in




FIG. 17.— Expanded plot of the spectrum of S-K2 of the IC 4040 EIG. A broad bump at around $\lambda 4650$ Å (Wolf-Rayet feature) can clearly be seen (the significance of the detection is $\sim 3\sigma$). The insertion is an enlarged spectrum around the Wolf-Rayet feature. Although signal-to-noise ratio is not sufficiently high to allow us to decompose the feature into individual emission lines, HeII $\lambda 4686$ Å and NIII $\lambda 4640$ Å can be identified.

particular, the [N II] line, of this knot.

4. DISCUSSION

4.1. Comparison with ram pressure stripping models

The one-sidedness of the morphologies of the EIGs of the Coma cluster strongly suggests RPS as the primary formation mechanism of these features (YY08; YY10). All four of the observed EIGs exhibit a monotonically-increasing pattern in their velocity fields. These types of velocity structures have been suggested in many numerical simulations of RPS (Vollmer & Huchtmeier 2003; Roediger & Hensler 2005; Roediger & Brüggen 2008; Tonnesen & Bryan 2009, 2010).

Recently, Tonnesen & Bryan (2009, 2010) performed three-dimensional hydrodynamical simulations of RPS incorporating radiative cooling. They found that radiative cooling fragments the disk gas of ram-pressured galaxy. This fragmentation forms holes in the disk and the holes accelerate the gas stripping. As a result, stripped gas flows straight behind the disk, such that the width of the wake of the stripped gas becomes much narrower than the width observed in models without cooling (Tonnesen & Bryan 2009). Tonnesen & Bryan (2010) examined observable values such as the density,




FIG. 22.— The velocity field of the EIG of IC 4040 (red circles) overlaid on that predicted by a numerical simulation made by Tonnesen & Bryan (2010). The observational data are not corrected for projection effect. The gray scaled contours represent the velocity field of the stripped gas in the simulation at a time 250 Myr after the collision between the model galaxy and the ICM (Tonnesen & Bryan 2010, reproduced by permission of the AAS). The collision velocity is 1413 km s^{-1} . The cut off temperature of the cooling curve of the model is $T_{\min} = 8000 \text{ K}$.

is comparable to that of the main stream of the wake of the simulation; $\approx 6 \text{ km s}^{-1} \text{ kpc}^{-1}$. Additionally, the simulation predicts that part of the stripped gas near the parent galaxy should have a very high relative velocity of $\sim 1000 - 1400 \text{ km s}^{-1}$ (Figure 22). This feature corresponds to SE-F1 (LV) of IC 4040 (SE-F1(LV)).

The main difference between the simulations and the observed data is the width of the velocity distribution. Numerical simulations made so far have predicted wide velocity distribution ($\sim 1000 \text{ km s}^{-1}$) perpendicular to the stream direction. Tonnesen & Bryan (2010) predicted much narrower velocity distribution (700 km s^{-1}) by incorporating radiative cooling, but it is still larger than that seen in the observations. The velocity fields of the RB199 and IC 4040 EIGs are well-ordered except for SE-F1(LV). The width of the velocity distribution of the EIG of IC 4040 is about $300 - 400 \text{ km s}^{-1}$, which is a factor of two narrower than the simulations. This discrepancy may be partly due to insufficient spatial coverage and detection limit in our observations. We observed only the bright parts of the EIG filaments and could not follow the faint thin clouds, which may have much higher velocities than the bright dense clouds. Further, the spatial coverage of our observation was limited by slits. Thus, our observation did not cover the entire velocity field of the EIGs. A wide field, two dimensional spectroscopy would be needed for further quantitative comparison between observations and the simulations.

The morphology of the wake of the stripped gas also differed between the simulations and the observation. While many simulations predict a widely spread wake, most EIGs in the Coma cluster have narrow, straight morphologies; the most remarkable example is the EIG around D100 (Yagi et al. 2007). The EIGs around GMP 2923 also has a narrow morphology. The EIGs around GMP 3071 and RB199 are wider, but the widths do not exceed the size of the galaxies. Evaporation of outer thin component of the stripped gas or some con-

finement mechanisms may narrow the morphology of the EIGs.

4.2. Star formation in the EIG

Strong star formation activities can be seen in the bright knots in the EIGs of RB199 and IC 4040. YY08 suggested, based on B , $H\alpha$ and $GALEX$ UV images, that the bright knots in the EIG of RB199 are star-forming regions. Smith et al. (2010) found asymmetric UV emission for 13 galaxies in the Coma center. They suggested that these emission regions were produced by active star formation in the stripped gas from their host galaxies. The sample of Smith et al. (2010) largely overlaps the EIG sample presented by YY10. Although YY10 did not identify optical continuum counterparts for most of the EIGs, Smith et al. (2010) pointed out that star formation activities are often associated with the EIGs (on the basis of their UV observations).

Recent numerical simulations of RPS have predicted active star formation in the stripped gas (Kronberger et al. 2008; Kapferer et al. 2008, 2009; Tonnesen & Bryan 2010; Tonnesen et al. 2011). YY08 reported good morphological agreement between the bright knots of the EIG of RB199 and the numerical simulation of Kronberger et al. (2008) (see also Kapferer et al. (2009)). Although the spatial resolution of these simulations is insufficient to reproduce molecular gas formation in the stripped gas, Yamagami & Fujita (2011) recently found that molecular clouds can be condensed in the RPS tails. They suggested that magnetic fields would play an important role in compressing the molecular clouds (Yamagami & Fujita 2011).




FIG. 23.— A $B - R_C$ vs. $FUV - B$ color-color diagram of the bright knots of the EIG of RB199 and IC 4040. Green points and orange points represent the data of the RB199 knots and the IC 4040 knots, respectively. Red line and blue dashed line show loci of star formation models (using PEGASE.2 (Fioc & Rocca-Volmerange 1997), see the text). Both models assume that stars are formed by infalling gas, and the infall rate is assumed to be proportional to $\exp(-t/t_i)/t_i$, where t_i is the timescale of the infall. We adopted $t_i = 100 \text{ Myr}$. Red line is the result of an exponentially decaying star formation model with a decay timescale of 100 Myr. Blue dashed line is the result of a constant star formation model.

We compared the broad-band colors of the star-forming knots of RB199 (knot 1 to knot 5) and IC 4040 (S-K2) with star formation model. Figure 23 is a $B - R_C$

versus $FUV - B$ color-color plot for the star-forming knots. The loci of star formation model calculations using PEGASE.2 (Fioc & Rocca-Volmerange 1997) are also plotted in the figure. In these calculations, we assumed that stars are formed by infalling of gas. The infall rate was assumed to be proportional to $\exp(-t/t_i)/t_i$, where t_i is the infall timescale. We adopted $t_i = 100$ Myr. We also assumed that the initial mass function (IMF) follows Salpeter IMF from $0.1 M_\odot$ to $120 M_\odot$, and the abundance of the infall gas is $0.6 Z_\odot$. We did not include emission-line components in color calculation. Figure 23 presents two cases; an exponentially-decaying star formation model with a decay timescale of 100 Myr, and a constant star formation model. YY08 showed similar plots for the RB199 knots. However, we found that the filter transmission functions of the *GALEX* UV bands adopted by YY08 were incorrect, and the predicted $FUV - B$ colors were significantly shifted to the blue direction. Additionally, they did not subtract $H\alpha$ emission contribution from the R_C band data. In this work, we used the correct *GALEX* FUV band transmission curve (Morrissey et al. 2005) to calculate FUV magnitude of the models, and the R_C magnitudes from which the $H\alpha$ emission flux was carefully subtracted based on our spectroscopy.

Figure 23 shows that the star-forming knots are very young, on the order of 100 Myr. These bright knots are located at ~ 20 kpc from the parent galaxies with a current relative velocity of $\sim 200 \text{ km s}^{-1}$. Thus it would take at least ~ 100 Myr for these knots to reach the current positions. This is comparable to or slightly shorter than the age estimation for the knots. However, the real travel time of the knots would be a few times longer than this value, because the knots are currently accelerated. Together with the low metal abundance nature of the knots (see section 3.2), this thus suggests that the most part of the stellar component of these knots was formed in the stripped metal poor gas which originally resided in the outskirts of the parent galaxies. Current active star formation associated with the knots also supports this idea. Using hybrid N-body / hydrodynamical simulations, Kapferer et al. (2009) found that 95% of stars newly formed by RPS reside in the wake of stripped gas. In their simulations, the timescale of star formation in the wakes is ~ 100 Myr (Kapferer et al. 2009). The young age of the star-forming knots of the EIGs suggested by our results is consistent with their findings.

4.3. Overlapped ionized gas in the EIG of RB199

We found that an ionized gas cloud overlaps the line of sight to $H\alpha$ cloud 2 of RB199. This cloud has a radial velocity of 6500 km s^{-1} , which is much slower than that of the EIG ($\sim 8000 \text{ km s}^{-1}$) and closer to the mean velocity of the Coma cluster. It indicates that this overlapped cloud is kinematically independent with the EIG.

What is this cloud and where did it originate? One possibility is that it is a gas cloud stripped from a nearby galaxy. A candidate of the parent galaxy is an early type galaxy (SDSS J125844.32+274251.6) near $H\alpha$ cloud 2. According to the SDSS database⁷, the color of this galaxy is that of a typical of S0 galaxies. Unfortunately, the recession velocity of this galaxy is not known. No

morphological indication suggests a physical connection between this galaxy and the overlapped cloud. We thus concluded that this galaxy is plausibly not the parent of the cloud.

On the other hand, this cloud may be a remnant of a stripping event which occurred long ago. In this case, identifying the parent galaxy would be difficult because an apparent physical connection between them would be almost completely lost.

Recently, a huge intra-cluster stellar population has been found in local- and intermediate-redshift clusters (Gregg & West 1998; Gonzalez et al. 2000; Mihos et al. 2005; Krick & Bernstein 2007). Such population is thought to be originated by galaxy-galaxy interaction or merger in the central region of the clusters. Tidal force associated with galaxy-galaxy interaction would strip the stars from the galaxies and spread them into the intra-cluster space. Associated with these tidal stripping events, dense disk gas may be also stripped, and a dense gas cloud may be floating in the intra-cluster space.

Although it is impossible to reach any conclusions from this rare example, this finding may suggest that there is a population of ‘floating ionized gas’ in clusters of galaxies (Gerhard et al. 2002; Cortese et al. 2004). Deep $H\alpha$ imaging of intra-cluster space and spectroscopic follow up will reveal how common this kind of objects are in galaxy clusters.

4.4. Fate of the EIG

Our spectroscopic observations showed that it is plausible that the EIGs associated to the four galaxies, RB199, IC 4040, GMP 2923 and GMP 3071, are formed by RPS. The observations also made it clear that strong star formation occurs in some parts of the stripped gas.

The EIGs around galaxies in the Coma cluster may act as a snapshot of a formation process of very faint galaxy population in clusters. The typical brightness and size of the bright knots in the EIGs are $M_R \sim -12 - -13$ and $1 - 2$ kpc (corresponding half light radius $\approx 200 - 300$ pc), respectively, which are similar to those of ultra compact dwarf galaxies (Drinkwater et al. 2004) or small dwarf spheroidals. This suggests that the bright knots are gravitationally self-bound systems⁸. In this case, they would survive and float in the halo of the parent galaxies or in the intra-cluster space of the Coma. After the star formation ceased, the luminosities of the bright knots will significantly decrease. Assuming the exponentially decaying star-formation model we used in Section 5.2, and assuming a typical age for the bright knots is 200 Myr, we can conclude that the luminosities of the knots would decrease by 2 – 3 magnitudes in R band in 1 Gyr. That is, the absolute magnitudes of the knots will eventually be $M_R \sim -10 - -11$. This luminosity range is within that of the dwarf spheroidal galaxies in the local group (Belokurov et al. 2007).

Most of the stars of the EIGs may be bound to the gravitational field of the parent galaxies. Kronberger et al. (2008) discussed that most of the stars formed in the stripped gas in their simulations are grav-

⁷ <http://www.mpa-garching.mpg.de/SDSS/DR7/>

⁸ Of course, measuring stellar velocity dispersion of the knots is necessary to check whether they are really self bound systems or not, but the spectral resolution and coverage of our spectroscopic data are not enough to perform such study.

itationally bound to the host galaxy and will fall back to the galaxy within 1 Gyr. In this case, the bright knots may be observed as features like ‘galaxy aggregates’ identified in the Coma cluster by Conselice & Gallagher (1998) in the fall-back process. Because stellar systems are collision-less ones, the stars will penetrate the galaxy disk and oscillate between two sides of the disk. Accordingly, the bright knots and the blue filaments will eventually be destroyed and distributed widely in the galaxy halo; in other words, the knots would form the halo population of the galaxy.

However, our spectroscopy revealed that the ionized gas of the knots has the relative velocity of $200 - 270 \text{ km s}^{-1}$ (Table 3). Because the stellar components of the knots are closely associated with the ionized gas, the stars would have almost the same velocity as the ionized gas. In this case, the knots probably escape from the gravitational potential of the parent galaxies and float into the intra-cluster space. In this context, it is interesting to note that an intra-cluster H II region was found in the Virgo cluster (Gerhard et al. 2002).

5. SUMMARY

We have presented the results of deep imaging and spectroscopic observations of very extended ionized gas (EIG) around four member galaxies of the Coma cluster, RB199, IC 4040, GMP 2923 and GMP 3071. Among them, the bright EIG around IC 4040 is remarkable and has complex morphology. It consists of two streams extending towards the southeast and southern directions from the galaxy. The southeast stream is connected to the central active star formation region. Many small ionized clouds are distributed between the southern stream and IC 4040. Several star-forming blue knots are embedded in the EIGs of RB199 and IC 4040. The extension of the EIGs ranges from $\sim 35 \text{ kpc}$ to $\sim 80 \text{ kpc}$.

The relative radial velocities of the EIGs with respect to the systemic velocity of the parent galaxies from which they emanate increase almost monotonically with the distance from the nucleus of the respective galaxies, reaching $\sim -400 - -800 \text{ km s}^{-1}$ at around $40 - 80 \text{ kpc}$ from the galaxies. The velocity decreasing rates are $-7 - -10 \text{ km s}^{-1} \text{ kpc}^{-1}$ for RB199, IC 4040 and GMP 2923. The GMP 3079 EIG has the steepest velocity gradient of $-25 \text{ km s}^{-1} \text{ kpc}^{-1}$. The kinematics of the EIGs is consistent with the concept that high-speed collisions between the intra-cluster medium and infalling galaxies of the Coma cluster form these features, because all the sample galax-

ies have systemic velocities much higher than the mean recession velocity of the cluster. We obtained a detailed velocity field for the bright EIG of IC 4040. The velocity dispersions of the diffuse filaments of the IC 4040 EIG reach $\sim 200 - 400 \text{ km s}^{-1}$. We also found a very low velocity filament at the southeastern edge of the disk of IC 4040. The filament has a velocity of -1300 km s^{-1} relative to the systemic velocity of the galaxy.

Some of the bright compact knots in the EIGs of RB199 and IC 4040 have blue continuum and strong H α emission. The equivalent widths of the H α emission of the knots range $200 - 1500 \text{ \AA}$. The emission line intensity ratios of the knots are basically consistent with those of sub-solar abundance H II regions. These facts indicate that intensive star formation occurs in the knots. Some filaments, including the low velocity filament of the IC 4040 EIG, exhibit shock-like emission line spectra, suggesting that shock heating plays an important role in ionization and excitation of the EIGs.

The morphologies and velocity fields of the EIGs are broadly consistent with those predicted by numerical simulations of ram pressure stripping. Our spectroscopic observations did not cover the entire regions of the EIGs due to spatial limitation of slits. This prevented us from making detailed comparison between the observation and the simulations. Deep and wide integral field spectroscopy would allow us to compare the kinematics of the EIGs with the prediction of numerical simulations in detail.

We are grateful to the staff of the Subaru telescope for their kind help with the observations. We thank the anonymous referee for valuable comments which significantly improve this paper. This work was in part carried out using the data obtained by a collaborative study on the Coma cluster. This study was in part carried out using the facilities at the Astronomical Data Analysis Center (ADAC), National Astronomical Observatory of Japan. This research made use of NASA’s Astrophysics Data System Abstract Service, NASA/IPAC Extragalactic Database, SDSS skyscraper, and GALEX GR5 database. This work was financially supported in part by Grant-in-Aid for Scientific Research No.23244030 from the Japan Society for the Promotion of Science (JSPS) and No.19047003 from the Ministry of Education, Culture, Sports, Science and Technology (MEXT).

REFERENCES

- Abadi, M. G., Moore, B., & Bower, R. G. 1999, *MNRAS*, 308, 947
 Abramson, A., et al. 2011, *AJ*, 141, 164
 Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, *ARA&A*, 47, 481
 Bekki, K. & Couch, W. J. 2003, *ApJ*, 596, L13
 Belokurov, V., et al. 2007, *ApJ*, 654, 897
 Boselli, A., & Gavazzi, G. 2006, *PASP*, 118, 517
 Boselli, A., Boissier, S., Cortese, L., & Gavazzi, G. 2008, *ApJ*, 674, 742
 Bravo-Alfaro, H., Cayatte, V., van Gorkom, J. H., & Balkowski, C. 2000, *AJ*, 119, 580
 Bravo-Alfaro, H., Cayatte, V., van Gorkom, J. H., & Balkowski, C. 2001, *A&A*, 379, 347
 Butcher, H., & Oemler, A. 1978, *ApJ*, 219, 18
 Butcher, H., & Oemler, A. 1984, *ApJ*, 285, 426
 Byrd, G., & Valtonen, M. 1990, *ApJ*, 350, 89
 Cayatte, V., van Gorkom, J. H., Balkowski, C., & Kotanyi, C. 1990, *AJ*, 100, 604
 Cayatte, V., Kotanyi, C., Balkowski, C., & van Gorkom, J. H. 1994, *AJ*, 107, 1003
 Chemin, L., et al. 2005, *A&A*, 436, 469
 Chung, A., van Gorkom, J. H., Kenney, J. D. P., Crawl, H., & Vollmer, B. 2009, *ApJ*, 659, L115
 Chung, A., van Gorkom, J. H., Kenney, J. D. P., Crawl, H., & Vollmer, B. 2009, *ApJS*, 138, 1741
 Conselice, C. J., & Gallagher, J. S. 1998, *MNRAS*, 297, L34
 Cortese, L., Gavazzi, G., Boselli, A., & Iglesias-Paramo, J. 2004, *A&A*, 416, 119
 Cortese, L., et al. 2006, *A&A*, 453, 847
 Cortese, L., et al. 2007, *MNRAS*, 376, 157
 Cortese, L., et al. 2010, *A&A*, 518, L49

- Cortese, L., Catinella, B., Boissier, S., Boselli, A., & Heinis, S. 2011, *MNRAS*, 415, 1797
- Crowl, H. H., Kenney, J. D. P., van Gorkom, J. H., & Vollmer, B. 2005, *AJ*, 130, 65
- Dressler, A. 1980, *ApJ*, 236, 351
- Dressler, A. 1994, *ApJ*, 430, 107
- Drinkwater, M. J. et al., 2004, *PASA*, 21, 375
- Fioc, M., & Rocca-Volmerange, B. 1997, *A&A*, 326, 950
- Fujita, Y., & Nagashima, M. 1999, *ApJ*, 516, 619
- Fujita, Y. 2004, *PASJ*, 56, 29
- Fumagalli, M., Gavazzi, G., Scaramella, R., & Franzetti, P. 2011, *A&A*, 528, A46
- Gavazzi, G., et al. 2001, *ApJ*, 563, L24
- Gavazzi, G., et al. 2003, *ApJ*, 597, 210
- Gavazzi, G., Fumagalli, M., Cucciati, O., & Boselli, A. 2010, *A&A*, 517, A73
- Gerhard, O., Arnaboldi, M., Freeman, K. C., & Okamura, S. 2002, *ApJ*, 580, L121
- Gonzalez, A. H., Zabludoff, A. I., & Zaritsky, D. 2000, *ApJ*, 618, 195
- Gregg, M. D., & West, M. J. 1998, *Nature*, 396, 549
- Gunn, J. E., & Gott, J. R. 1972, *ApJ*, 176, 1
- Henriksen, M., & Byrd, G. 1996, *ApJ*, 459, 89
- Hester, J. A., et al. 2010, *ApJ*, 716, L14
- Ho, L. C., Filippenko, A. V., & Sargent, W. L. W. 1997, *ApJS*, 112, 391
- Jáchym, P. et al. 2007, *A&A*, 472, 5
- Jansen, R. A., Fabricant, D., Franx, M., & Caldwell, N. 2000, *ApJS*, 126, 331
- Kapferer, W., Kronberger, T., Ferrari, C., Riser, T., & Schindler, S. 2008, *MNRAS*, 389, 1405
- Kapferer, W., Sluka, C., Schindler, S., Ferrari, C., & Ziegler, B. 2009, *A&A*, 499, 87
- Kashikawa, N., et al. 2002, *PASJ*, 54, 819
- Kauffmann, G., et al. 2003, *MNRAS*, 346, 1055
- Kennicutt, R. C. 1998, *ApJ*, 498, 541
- Kenney, J. D. P., & Koopmann, R. A. 1999, *AJ*, 117, 181
- Kenney, J. D. P., Tal, T., Crowl, H. H., Feldmeier, J., & Jacoby, G. H. 2008, *ApJ*, 687, L69
- Kewley, L. J., Dopita, M. A., Sutherland, R. S., Heisler, C. A., & Trevena, J. 2001, *ApJ*, 556, 121
- Kewley, L. J., & Dopita, M. A. 2002, *ApJS*, 142, 35
- Kong, X., Cheng, F. Z., Weiss, A., & Charlot, S. 2002, *A&A*, 396, 503
- Koopmann, R., & Kenney, J. D. P. 2004, *ApJ*, 613, 866
- Krick, J. E., & Bernstein, R. A. 2006, *AJ*, 134, 466
- Kronberger, T., Kapferer, W., Ferrari, C., Unterguggenberger, S., & Schindler, S. 2008, *A&A*, 481, 337
- Mihos, J. C., Harding, P., Feldmeier, J., & Morrison, H. 2005, *ApJ*, 631, L41
- Miyazaki, S. et al. 2002, *PASJ*, 54, 833
- Moore, B., Katz, N., Lake, G., Dressler, A., & Oemler, A. 1996, *Nature*, 379, 613
- Morrissey, P. et al. 2005, *ApJ*, 619, L7
- Okamoto, T., & Nagashima, M. 2003, *ApJ*, 587, 500
- Okamura, S., et al. 2002, *PASJ*, 54, 883
- Oosterloo, T., & van Gorkom, J. 2005, *A&A*, 437, L19
- Postman, M., & Geller, M. J. 1984, *ApJ*, 281, 95
- Randall, S., et al. 2008, *ApJ*, 688, 208
- Roediger, E., & Hensler, G. 2005, *A&A*, 433, 875
- Roediger, E., Brüggen, M., & Hoeft, M. 2006, *MNRAS*, 371, 609
- Roediger, E., & Brüggen, M. 2007, *MNRAS*, 380, 1399
- Roediger, E., & Brüggen, M. 2008, *MNRAS*, 388, 465
- Sakai, S., et al. 2002, *ApJ*, 578, 842
- Schulz, S., & Struck, C. 2001, *MNRAS*, 328, 185
- Sivanandam, S., Rieke, M. J., & Rieke, G. H. 2010, *ApJ*, 717, 147
- Smith, G. P., Treu, T., Ellis, R. S., Moran, S. M., & Dressler, A. 2005, *ApJ*, 620, 78
- Smith, R. J., et al. 2010, *MNRAS*, 408, 1417
- Solanes, J. M., et al. 2001, *ApJ*, 548, 97
- Sun, M., Donahue, M., & Voit, M. 2007, *ApJ*, 671, 170
- Sun, M., et al. 2010, *ApJ*, 708, 946
- Tonnesen, S., & Bryan, G. L. 2009, *ApJ*, 694, 789
- Tonnesen, S., & Bryan, G. L. 2010, *ApJ*, 709, 1203
- Tonnesen, S., Bryan, G. L., & Chen, R. 2011, *arXiv:1103.3273*
- Trentham, N., Sampson, L., & Banerji, M. 2005, *MNRAS*, 357, 783
- van der Wel, A., et al. 2007, *ApJ*, 670, 206
- van der Wel, A., Bell, E. F., Holden, B. P., Skibba, R. A., & Rix, H.-W. 2010, *ApJ*, 714, 1779
- van Gorkom, J. H. 2004, in *Clusters of Galaxies: Probes of Cosmological Structure and Galaxy Evolution*, ed. J. S. Mulchaey, A. Dressler, & A. Oemler (Cambridge: Cambridge Univ. Press), 305
- Veilleux, S., & Osterbrock, D. M. 1987, *ApJS*, 63, 295
- Vollmer, B. 2009, *A&A*, 502, 427
- Vollmer, B., Braine, J., Balkowski, C., Cayatte, V., & Duschl, W. J. 2001, *A&A*, 374, 824
- Vollmer, B., Cayatte, V., Balkowski, C., & Duschl, W. J. 2001, *ApJ*, 561, 708
- Vollmer, B., & Huchtmeier, W. 2003, *A&A*, 406, 427
- Vollmer, B., & Huchtmeier, W. 2007, *A&A*, 462, 93
- Vollmer, B., Brane, J., Pappalardo, C., & Hily-Blant, P. 2008, *A&A*, 491, 445
- Vollmer, B., et al. 2009, *A&A*, 496, 669
- Yamagami, T., & Fujita, Y. 2011, *PASJ*, 63, 1165
- Yagi, M., et al. 2007, *ApJ*, 660, 1209
- Yagi, M., et al. 2010, *AJ*, 140, 1814 (YY10)
- Yoshida, M., et al. 2002, *ApJ*, 567, 118
- Yoshida, M., et al. 2004, *AJ*, 127, 90
- Yoshida, M., et al. 2008, *ApJ*, 688, 918 (YY08)